

# Deccox 10X

## Suthers

### Medicated Type B

361

Mixing Mineral for Cattle in Confinement

For the prevention of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii*.

#### Active Drug Ingredient

Decoquinatate.....4540.0 g/ton

(0.4994%)

#### Guaranteed Analysis

Crude Protein			Min.	16.0%
Crude Fat			Min.	6.4
Crude Fiber			Min.	5.2
Calcium	Min	3.8	Max.	4.8
Potassium			Min.	0.9%

#### Ingredients

Corn Dried Distillers Grains with Solubles, Magnesium Mica, Calcium Carbonate, Vegetable Oil.

#### Feeding Directions:

Mix this Type B medicated feed with non-medicated feed ingredients to manufacture one ton of complete feed containing 12.9-90.8 g/ton decoquinatate. The resulting Type C medicated feed should be fed for at least 28 days during periods of exposure to coccidiosis or when it is likely to be a hazard. The concentration of decoquinatate in the final Type C feed will vary with body weight and feed consumption to provide 22.7 mg per 100 pounds of body weight per day.

Refer to the table below for mixing directions.

Decoquinatate Level Desired g/ton	Type B to Mix lb/ton	Non-Med to mix Lb/ton
12.9	5.68	1994.32
22.7	10.00	1990.00
50	22.03	1977.97
75	33.04	1966.96

**Warning:** Do not feed to cows producing milk for food. Feed for at least 28 days during periods of coccidiosis exposure or when experience indicates coccidiosis to be a hazard. Feed Type C feed or milk replacer to provide 22.7 mg per 100lb. bodyweight (0.5 mg/kg) per day.

**Caution: Do not use in feeds containing bentonite.**