Apex[™] Plus for Dogs

Ony for Dogs & Puppies 5-22lbs & over 8 weeks of Age

3-month Supply

(Contains fipronil and (S)- Methoprene, the active ingredients used in FRONTLINE® Plus for Dogs. *)

Kills Fleas, Flea Eggs, Flea Larvae, Ticks & Chewing Lice

Waterproof 24 hours after application

Kills ticks, including those that may transmit Lyme disease.

Active Ingredients:

Fipronil	9.80%
(S)-Methoprene	8.80%
Other Ingredients	81.40%
Total	100.00%

^{*}This product is not manufactured or distributed by Boehringer Ingelheim Animal Health USA, Inc., the makers of FRONTLINE® Plus for Dogs. FRONTLINE® is a registered trademark of Boehringer Ingelheim Animal Health USA, Inc.

Read entire label before each use. Use only on dogs & puppies over 8 weeks of age and 5-22lbs.

Keep out of reach of Children

Precautionary Statements

Hazards to Humans. Caution.

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with skin., eyes or clothing. Wash hands thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

Hazards to Domestic Animals

For external Use on dogs only. Individual sensitivities, while rare, may occur after using any pesticide product on your dog. Certain medications can interact with pesticides. Do not treat your dog with more than one pesticide product at a time. Overdosing your dog can result in serious illness or even death. Consult a veterinarian before using on medicated, debilitated or aged pregnant or nursing dogs. Consult a veterinarian before using on dogs with known organ disfunction. If your dog is exhibiting signs of and /or is being treated for skin dermatitis, talk to your veterinarian before applying any topical flea and tick product. This product is not a substitute for prescribed heartworm preventative.

First Aid

IF SWALLOWED: Call a poison control center or a doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or a doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

Hotline Number: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at 1-866-897-8050 for emergency medical treatment information.

Directions for Use

It is in violation of the Federal law to use this product in a manner inconsistent with its labeling.

Do NOT allow children to apply product.

To prevent harm to you and your dog, read entire label before each use. Follow all directions and precautionary statements carefully.

Restrictions:

Do not use on cats, rabbits or animals other than dogs. Do not allow your dog to ingest this product. Do not use on dogs or puppies under 8 weeks of age or less than 5lbs. Use entire contents of applicator on each dog. Weigh your dog to be sure you are applying the right dose formulated for the weight of your dog. Separate the treated dog from all other cats and dogs for 24 hours after treatment has been applied. Do not apply more than one applicator per treatment for one dog regardless of dog's weight. Do not bathe or allow water immersion (swimming) for 24 hours after application. The dog needs to be dry at the time of the application and remain dry for 24 hours after the application. Do not apply more often than once every 30 days.

How to Open:

- 1. Tear through perforation
- 2. Cut with scissors to open applicator

How to apply:

- 1. Invert the applicator over the dog and use open end to part dog's hair
- 2. Squeeze the applicator firmly to apply all of the solution to the dog's skin from the back of the neck to a point midway between the neck and tail.

Frequency of Application:

Use APEX™ Plus for dogs every 30 days for control of flea, tick and chewing lice infestations. Apply monthly to prevent flea infestation to your cat. Apply every 30 days to control ticks and chewing lice. This product remains effective, even after bathing, water immersion, or exposure to sunlight 24 hours after application. Allow treated area to dry thoroughly before touching. Dog should be dry at time of application and for 24 hours after application. Do not reapply for 30 days.

Side Effects:

Monitor your dog after application. Dogs may experience temporary startle effects when any product is applied. Cats may experience some temporary irritation at the site of product application such as redness, scratching or other signs of discomfort. If signs of sensitivity occur, bathe your dog with mild soap and rinse with large amounts of water. Gastrointestinal signs such as vomiting or diarrhea have also been reported. If these or other side effects occur consult your veterinarian or call 1-866-897-8050. If your cat has an unusual reaction to the initial application, consult a veterinarian before repeating application.

Do not Use on Cats: Keep cats away from treated dogs for 24 hours. If applied to a cat, contact your veterinarian.

Storage and Disposal

Pesticide Storage: Do not remove applicator from the pack until ready to use. Store in a cool (below 77°F) dry place inaccessible to children and pets. Do not refrigerate. Protect from direct sunlight.

Pesticide Disposal and Container Handling: Non-refillable container. Do not reuse or refill this container. IF EMPTY: Offer recycling if available. IF PARTLY FILLED: Call your local solid waste agency for disposal instructions. Never place unused product sown any indoor or outdoor drain.